

ISSN 2518-1467 (Online),
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Ш Ы С Ы

ВЕСТНИК

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

THE BULLETIN

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

1944 ЖЫЛДАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С 1944 ГОДА
PUBLISHED SINCE 1944

4

АЛМАТЫ
АЛМАТЫ
ALMATY

2018

JULY
ИЮЛЬ
ШІЛДЕ

NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

М. Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

Абиев Р.Ш. проф. (Ресей)
Абишев М.Е. проф., корр.-мүшесі (Қазақстан)
Аврамов К.В. проф. (Украина)
Аппель Юрген проф. (Германия)
Баймуқанов Д.А. проф., корр.-мүшесі (Қазақстан)
Байпақов К.М. проф., академик (Қазақстан)
Байтулин И.О. проф., академик (Қазақстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Қазақстан)
Велихов Е.П. проф., РҒА академигі (Ресей)
Гашимзаде Ф. проф., академик (Әзірбайжан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., корр.-мүшесі (Қазақстан)
Джрбашян Р.Т. проф., академик (Армения)
Қалимолдаев М.Н. проф., академик (Қазақстан), бас ред. орынбасары
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., корр.-мүшесі (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалықов Ж.У. проф., академик (Қазақстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., корр.-мүшесі (Қазақстан)
Полещук О.Х. проф. (Ресей)
Поняев А.И. проф. (Ресей)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Қазақстан)
Таткеева Г.Г. проф., корр.-мүшесі (Қазақстан)
Умбетаев И. проф., академик (Қазақстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., РҒА корр.-мүшесі (Ресей)
Якубова М.М. проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
www: nauka-nanrk.kz, bulletin-science.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. х. н., проф. академик НАН РК
М. Ж. Журинов

Р е д а к ц и о н н а я к о л л е г и я:

Абиев Р.Ш. проф. (Россия)
Абишев М.Е. проф., член-корр. (Казахстан)
Аврамов К.В. проф. (Украина)
Апель Юрген проф. (Германия)
Баймуканов Д.А. проф., чл.-корр. (Казахстан)
Байпаков К.М. проф., академик (Казахстан)
Байтулин И.О. проф., академик (Казахстан)
Банас Иозеф проф. (Польша)
Берсимбаев Р.И. проф., академик (Казахстан)
Велихов Е.П. проф., академик РАН (Россия)
Гашимзаде Ф. проф., академик (Азербайджан)
Гончарук В.В. проф., академик (Украина)
Давлетов А.Е. проф., чл.-корр. (Казахстан)
Джрбашян Р.Т. проф., академик (Армения)
Калимолдаев М.Н. академик (Казахстан), зам. гл. ред.
Лаверов Н.П. проф., академик РАН (Россия)
Лупашку Ф. проф., чл.-корр. (Молдова)
Мохд Хасан Селамат проф. (Малайзия)
Мырхалыков Ж.У. проф., академик (Казахстан)
Новак Изабелла проф. (Польша)
Огарь Н.П. проф., чл.-корр. (Казахстан)
Полещук О.Х. проф. (Россия)
Поняев А.И. проф. (Россия)
Сагиян А.С. проф., академик (Армения)
Сатубалдин С.С. проф., академик (Казахстан)
Таткеева Г.Г. проф., чл.-корр. (Казахстан)
Умбетаев И. проф., академик (Казахстан)
Хрипунов Г.С. проф. (Украина)
Юлдашбаев Ю.А. проф., член-корр. РАН (Россия)
Якубова М.М. проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

© Национальная академия наук Республики Казахстан, 2018

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

M. Zh. Zhurinov

E d i t o r i a l b o a r d:

Abiyev R.Sh. prof. (Russia)
Abishev M.Ye. prof., corr. member. (Kazakhstan)
Avramov K.V. prof. (Ukraine)
Appel Jurgen, prof. (Germany)
Baimukanov D.A. prof., corr. member. (Kazakhstan)
Baipakov K.M. prof., academician (Kazakhstan)
Baitullin I.O. prof., academician (Kazakhstan)
Joseph Banas, prof. (Poland)
Bersimbayev R.I. prof., academician (Kazakhstan)
Velikhov Ye.P. prof., academician of RAS (Russia)
Gashimzade F. prof., academician (Azerbaijan)
Goncharuk V.V. prof., academician (Ukraine)
Davletov A.Ye. prof., corr. member. (Kazakhstan)
Dzhrbashian R.T. prof., academician (Armenia)
Kalimoldayev M.N. prof., academician (Kazakhstan), deputy editor in chief
Laverov N.P. prof., academician of RAS (Russia)
Lupashku F. prof., corr. member. (Moldova)
Mohd Hassan Selamat, prof. (Malaysia)
Myrkhalykov Zh.U. prof., academician (Kazakhstan)
Nowak Isabella, prof. (Poland)
Ogar N.P. prof., corr. member. (Kazakhstan)
Poleshchuk O.Kh. prof. (Russia)
Ponyaev A.I. prof. (Russia)
Sagiyani A.S. prof., academician (Armenia)
Satubaldin S.S. prof., academician (Kazakhstan)
Tatkeyeva G.G. prof., corr. member. (Kazakhstan)
Umbetayev I. prof., academician (Kazakhstan)
Khripunov G.S. prof. (Ukraine)
Yuldashbayev Y.A., prof. corresponding member of RAS (Russia)
Yakubova M.M. prof., academician (Tadjikistan)

Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2518-1467 (Online),

ISSN 1991-3494 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

I. I. Mardanov¹, S. A. Tarikhazer², I. Y. Kuchinskaya², S. Y. Guliyeva², L. A. Ismaylova³

¹Sumgait State University, Sumgait, Azerbaijan,

²Institute of Geography named by acad. H. A. Aliyev of ANAS, Baku, Azerbaijan,

³Azerbaijan State Oil and Industry University, Baku, Azerbaijan.

E-mail: geography.sumqayit@mail.ru, kerimov17@gmail.com, Irina.danula@gmail.com,
selinsevil@gmail.com, Latifa.isamylova@gmail.com

GEOSYSTEM ANALYSIS OF DEVELOPMENT OF LANDSLIDE LANDSCAPES OF THE SOUTH-EASTERN SLOPE OF GREATER CAUCASUS

Abstract. The southeast tip of Greater Caucasus which is entering borders of Azerbaijan and being part of the Alpine and Himalaya geosynclinal belt, is characterized by high dynamism exogenous the relief forming the processes having essential impact on development of a landscape situation. In this regard there is a need of development of various methods for creation of scenarios of possible changes of a geoecological situation in various massif having harmful consequences, by identification of relationships of cause and effect. For this purpose, in this work possibilities of forecasting of landslides in natural area of Greater Caucasus with use of available geological and geomorphological, climatic and landscape data, and also visual supervision were analysed. These data allowed to reveal the main distinctions of factors of a relief situation, hydroweather conditions in certain physiographic areas in which landslides, character and intensity of the land use, to some extent influencing a descent of landslides and subject to their destructive influence are shown.

Keywords: processes, slope, exodynamic, landscapes, erosion, factor, sub district, Caucasus.

1. Introduction. Questions of identification of the reasons of geodynamic processes – taluses, scatterings and landslides are many years in the center of attention of the geographical public of the country because of harmful consequences of these phenomena for the social sphere and economic activity of all mountain regions [14].

The southern and Northeast slope of Greater Caucasus differ the complex geomorphological structure, being shown in difficult structure of alternation of water-permeable layers of breeds, an interlacing of tectonic violations, and also the high seismicity which quite often is hardly noticeable, but a decisive factor of slipping and a collapse of mass of breeds.

The inclination of a terrestrial surface saving in considerable energy of all mass of breeds of a slope has essential impact on development of landslide processes. But, as we know, big biases of slopes not always lead to landslides, even on sites close located to landslide massifs. So, slopes with strong maternal breeds are steady, slopes with alternation of layers of friable breeds and clays are the most subject to influence of geodynamic factors. Big biases of slopes, especially characteristic for the Southern slope of Greater Caucasus Range lead to landslides of landslide character, in a root changing shape of a landscape of this site [12, 13].

On degree of stability to impact of landslide processes of the territory of a massif active and active sites are differentiated on rather steady, so-so. Within a mountain meadow zone steady sites in the landslide relation are characterized by existence of clearly expressed subalpine and Alpine landscapes. Such sites can be subject to influence of others exodynamic processes – to a soil erosion, accumulation and movement of taluses and the scatterings constituting not smaller danger to a landscape and all ecological situation (figure 1).




Figure 1 – Space image of part of the studied territory of a river basin of Girdymanchay of the 2010 with the allocated contours of the soils, different degree of erodibility. Scale 1 : 25 000;
1 - not bald-headed sites; 2 - slightly eroded sites; 3 - moderately eroded sites; 4 - severely eroded sites

It is known that an important element of development of landscapes is transfer of chemical particles as part of circulation of substances. In this regard mountain landscapes of Greater Caucasus are low-studied and demand the analysis of impact of exogenous processes on changes of quantity and a ratio of various chemical elements and their connections in soils and vegetation depending on intensity of various natural phenomena. For these purposes experts of various profile – geographers, soil scientists, botanists, the chemists which common efforts can help to reveal an overall picture of occurring geochemical changes can be attracted [17-19].

The geodynamic situation in Apsheron peninsula substantially becomes complicated the intensive town planning, being accompanied development of social and industrial infrastructure that leads to change of an initial relief and strengthening of factors of an landslide forming.

2. Objects and methods of researches. The southeast tip of Greater Caucasus which is active from the point of view of a descent of landslides, always drew attention of researchers of various disciplines – geologists, geomorphologists, landscape scientists, soil scientists, etc. trying to establish the reasons of this destructive process, to give an assessment of extent of influence of this or that factor on this phenomenon. The main landslide massifs, lithologic structure of breeds of landslide slopes, the basic landscape elements of separate large landslides were during this time defined [1-3].

These works were performed during field visual researches, and also the cameral researches meaning measurements on topographic maps, revealing the biases of a surface stimulating slipping of mountain masses.

Measurements of the cracks formed as a result of landslides on different massifs and definitions of their temporary dynamics allow to carry out in a certain degree territory division into districts on activity of breeds for the purpose of identification most of the landslide territories. The description of a landscape situation, its separate components promotes detection of the distinctions which were showed through a certain time occurring also under the influence of geodynamic processes.

The interpretation of space images gives the chance of them to interpretation for detection of geomorphological and landscape features of consequences of landslides both their spatial and temporary dynamics in the presence of pictures of different years. In Azerbaijan many years were spent works on creation and improvement of various means and methods of interpretation of space data for definition of the directions of development of the harmful natural phenomena [10-12].

3. The received results. Materials of numerous researches of an environment of Greater Caucasus available in existence can give the chance to reveal prerequisites of emergence and development of landslide processes and to create a certain group of landslide massifs for the purpose of development of protective measures on prevention of their harmful consequences. The material on physiographic and synoptic division into districts of natural area of Greater Caucasus within Azerbaijan was for this purpose processed and a number of the factors causing a descent of landslide masses is defined.

Divaryan landslide stream, beginning on a northwest slope of the mountain Matur, on the left coast of Girdymanchay, at the absolute height of 2200 m, it is stretched at distance of 2,9 km to the course of Girdymanchay and finishes the activity at the absolute height of 1400 m. In an amphitheater and a transitional zone width of a landslide stream reaches 10-15 m, and on a carrying out cone more than 2,5 km. Its landslide materials narrowed the course of Girdymanchay to 3-5 m. The modern landscape of a landslide difficult also is characterized by existence of cracks, ridges, shaft, bushes, hollows, pools, boggy sites and fresh landslide materials.

As the digital model of a relief of the Divaryan landslide, created on the basis of the GIS technology shows, wavy character of a surface causes diversity of landscape structure of landslide weight. Namely, camber of a middle part of a landslide with the best conditions of moisture accumulation, most likely, is the reason of development of shrubby and wood vegetation on relatively weak inclined sites, including, sites of a cone of carrying out (figure 2).




Figure 2 – Digital model of a relief of the Divaryan landslide located on the left river bank Girdymanchay. This model is received on the basis of processing of a space picture of high resolution of 2012 and 2013. Continuous horizontals are carried out through each 20 meters

The similar model is created and for the Demirchi landslide which has been also located on the Southeast slope of Greater Caucasus Range, which development down can influence changes of the bed of the river (figure 3). Unlike the Divaryan landslide the surface of this landslide has a hollow form. Demirchi landslide stream, beginning to the East from the Lagich pass in river Demirchichay headwaters (a river basin Pirsaat) at the absolute height of 2000 m, proceeds to 1700 m. Its length is 25 km, width is 100-500 m. It develops within mountain meadow and mountain-steppe landscape types.




Figure 3 – Digital model of a relief of the Demirchi landslide located in a river basin Pirsaat.
This model is received on the basis of processing of a space picture of high resolution of 2012 and 2013.
Continuous horizontals are carried out through each 20 meters

Mountain territories of Greater Caucasus are included in limits of several physiographic areas. The Gonagkend area covers in the structural relation the Tufan anticlinorium, the Shakhdag-Hyzisynclorium, the Tengi-Beshbarmag anticlinorium, east suburb of the Zagatalo-Govdagsynclorium and Hussarmonocline. Due to the formation at various absolute heights of a number of intermountain hollows conditions for development of settlements, agriculture and cattle breeding which however, can serve as the reason erosive, but not landslide process are created. The landslide phenomena here have the natural character connected, apparently, with fluctuations of seismic activity, nature of the spreading breeds, washing-away activity of the rivers, especially, during high waters and an atmospheric precipitation, including, storm rains. The intermountain hollow of Shakhdyuzu (2400-2700 m) is used as a summer pasture. The relief of the area possesses the big range of heights (200-4460 m) and therefore altitudinal zonality is presented here in a full range. Due to the fall in the southeast direction of absolute height of a relief and aridity strengthening in hillsides reduction of a river drain and disappearance of mountain landscape belts, an aridization of the woods, expansion of areas of dry steppes and semi-deserts towards low-mountainous is observed.

Height differences on the Southern slope of Greater Caucasus Range which generally covers the Zagatalo-Lagich physio geographic area which is stretching from the West, from the Georgian border to a river Girdimanchay valley in the east on 220 kilometers, with cool slopes are made by 2800-2900 meters. Biases of a surface change in limits 30°-45° that probably is a leading factor of development of landslide processes, along with showers, characteristic and for the Northeast slope of Greater Caucasus and high seismicity. Number of the rivers (Mazymchay, Belokanchay, Kurmukhchay, Dashagilchay, etc.) crosses the Southern slope across, forming valleys with cool slopes. In the territory the broad-leaved woods which however can't serve fixing from breed sliding by a factor, mountain meadows and subnival landscapes dominate.

Shemakhi (Mountain Shirvan) the physio geographic area surrounded from the West Akhsu's river, from the North a watershed of Greater Caucasus Range, from the East the Gobustan low-mountainous, and from the South the Shirvan steppe is characterized by high seismicity (8-9 points) which probably plays the leading role in coupling violation between the mass of breeds and a descent of landslides. In the territory of the area where unlike previous, with more humid climate, mountain-steppe, forest-steppe, mountain and forest and mountain meadow landscape complexes prevail.

Studying of a geodynamic situation in Apsheron which is part of Greater Caucasus becomes an actual scientific and practical problem for life support of the most part of the population of the country which has concentrated on the territory of the Baku agglomeration, enduring construction boom which is capable to aggravate earlier existing problems and to create new in connection with intensive high-rise construction in the Baku amphitheater. Today the main landslide massifs in city boundaries are known and the most probable causes of a descent of landslides on these sites are established. However with expansion of city boundaries and advance of housing construction, including, elite, for example, to the South from the downtown towards the Bail ledge there are new dangers to life of people on once weak-populated site of Apsheron.

The territory of Apsheron and the city of Baku is included in limits of the Gobustan-Apsheron physio geographic region of natural area of the Southeast tip of Greater Caucasus which is the landslide dangerous territory for all South Caucasus [4]. The most part of the area consists of low-mountainous with a wide circulation arid denudation processes and in the tectonic relation enters limits of the Shamakhi-Gobustan synclinorium and a southeast extremity of the mega anticlinorium of Greater Caucasus [11]. Owing to a relief of low-mountainous and plains, and also high aridity of climate in the territory landscapes of semi-deserts and dry steppes dominate. Here the halophytic plants is most of all developed.

The territory of Greater Caucasus is included in limits of several synoptic areas.

The Oguz-Ismailli area is exposed to influence of the centers of the cold air located over the Kara Sea, Scandinavia, the South of Eastern Europe and Kazakhstan, getting in the territory through the Caspian Sea and the Azor maximum. Seldom air masses get through the territory of Georgia. Sometimes warm air of a subtropical anti-cyclone gets. Three subdistricts are allocated.

In a high-mountain subdistrict average annual temperatures make 0° - 7°C , middle January temperature -5° - 10°C (frost), and middle July $+5^{\circ}$ - 17°C , an average annual amount of precipitation of 1000-1300 mm. The most damp period – the end of spring beginning of summer. During this period there can be an activization of a role of ground waters in development of landslides.

The Guba-Shamakhi area more than other areas is affected by cold air masses Arctic and middle latitudes and the Central Asian anti-cyclone. On the contrary, influence of a tropical anti-cyclone is shown here poorly, and the Azor maximum in comparison with other areas – is strong.

Mountain subdistrict – average annual temperature reaches $+5^{\circ}$ - 7°C , middle January $+4^{\circ}$ - 6°C , middle July $+14^{\circ}$ - 15°C less. The average annual amount of precipitation makes 400-600 mm that says that the role of an amount of precipitation in an landslide forming is less, than their quality in comparison with the previous area though the maximum of a precipitation here is observed, as well as in the Oguz-Ismailli area at the end of spring beginning of summer.

The territory of the Apsheron peninsula and the city of Baku is included in limits of the Apsheron-Gobustan synoptic area differing from others by strong North southern air streams. Influence of area of a high pressure formed Kara, Scandinavian and Azor maximum over Southeast Europe is the main reasons of it. Within this area two subdistricts are allocated: subdistrict of Apsheron and Gobustan subdistrict.

In a subdistrict of Apsheron average annual temperature makes $+14^{\circ}$ - 15°C , middle January $+3^{\circ}$ - 4°C , middle July $+24^{\circ}$ - 26°C , the average annual sum of a precipitation of 100-250 mm, the greatest number of a precipitation drops out in the fall, the winds of the northern direction which are called “xazri” (or the Baku north), often being accompanied a storm dominate. Storms, fogs and phenomenon snowfalls seldom repeating.

In the Gobustan subdistrict average annual temperature makes $+11^{\circ}$ - 13°C , middle January -1° - 3°C , middle July $+22^{\circ}$ - 25°C , the average annual sum of a precipitation of 150-300 mm, the greatest number of a precipitation drops out in the fall. Northern and northeast winds dominate, during the hair dryer the strong southwest wind blows. Storms, fogs and snowfalls are observed more often than in the Apsheron subdistrict.

Apparently from the provided data, climatic conditions of these subdistricts strongly don't differ and can't be considered as a leading factor of emergence of landslides.

4. Conclusions. The numerous researches directed on development of actions for prevention of a descent of landslides are based on the purposes of reduction of biases of hillsides, the phytomeliorative works meaning restoration of a close vegetable cover on broken massifs. Such approach proceeds from insufficient understanding of an essence of the mechanism of this process covering not simply terrestrial surface, but rather powerful cover of sedimentary breeds with a certain stratification, sometimes some tens meters. Quite often in publications of various character expression “sliding of soils” that proceeds from misunderstanding of a geological and geomorphological essence of landslide process, the superficial relation to possible consequences of this phenomenon meets.

At the same time, attempts of melioration of landslide massifs which are ineffective are made. They, are generally directed on alignment of a surface and restoration of a close soil and vegetable cover, including, by implementation a forest plantation.

These measures lead to temporary stabilization of a geodynamic situation that actually, is the beginning of a new cycle in development of landslide process. Planting of trees improves soil and ecological, but not a geological and geomorphological situation as roots of trees aren't capable to constrain movement of mass of breeds sufficiently. Numerous landslides within a mountain and forest belt of Greater Caucasus within Azerbaijan, strongly changed all shape of a landscape can serve as an example, introducing before not meeting elements in its horizontal structure.

5. Alleged actions. In this regard there is an urgent need in the large-scale inventory of landslide massifs meaning drawing up detailed landscape, geological and geomorphological, soil and geobotanical sketch maps, including, with use of materials of space shooting [23-25]. Cartographic materials have to be added with the meteorological data including information on an annual course of air temperature, quantity, character and a mode of loss of a precipitation, average annual and average monthly air temperatures.

In the description of the landslide massif information on a humanitarian and economic situation of the landslide site, including data on number of settlements, number of their population, type of the settlement, features of their arrangement, quantity and character of the enterprises, the social objects located in this territory has to take an important place.

Accumulation and processing of large volume of data is able to afford to carry out comparison of the probable factors causing process of a descent of landslides and to reveal the key factor which hasn't been considered by initial consideration, for the purpose of search of opportunities of its neutralization [9, 12, 13, 20-22].

REFERENCES

[1] Alizadeh E.K., Guliyeva S.Y., Tarikhazer S.A. (2005) The assesment of degree of susceptibility of geocomplexes of the Southern slope of Greater Caucasus landslide processes // In Scientific and practical conference Natural and Destructive Phenomena of Sheki-Zakatala Area and Ecogeographical Problems of Development of the Region, on June 9-10. Sheki: 63-65. (in Russian).

[2] Alizadeh E.K., Guliyeva S.Y., Kuchinskaya I.Y., Tarikhazer S.A. (2015) Intensity of landslips in mountain geosystems of Azerbaijan and its estimation // International Journal of Scientific Research and Innovative Technology. ISSN 2313-3759. Vol. 2, 10, October. (in Eng.).

[3] Alizadeh E., Ismayilov M., Guliyeva S., Zeynalova S., Tarikhazer S., Yunusov M., Mustafayev N., Mammadbayov E. (2016) The assessment of landscape and environmental risks and hazards caused by landslides in mountain areas. Applied ecology and environmental research, 14(3), ALÖKI Kft., Budapest, Hungary: 573-586. (in Eng.).

[4] Budagov B.A., Alizadeh E.K., Tarikhazer S.A. (2005) Current trends of development it is spontaneous – destructive processes and an assessment of ecogeomorphological danger (on the example of the southern slope of Greater Caucasus) Inthe Scientific and practical conference Natural and Destructive Phenomena of Sheki-Zakatala Area and Ecogeographical Problems of Development of the Region, on June 9-10. Sheki, 2005: 25-28. (in Russian).

[5] Etzelmuller B., Romstad B., Fjellanger J. (2007) Automatic regional classification of topography in Norway // Norwegian Journal of Geology. Vol. 87: 167-180. (in Eng.).

[6] Garibov Y., Mardanov I., Ismaylova N., Ahmadova G., Aliyeva R. (2016) Ekzogenetic features landscapes of high mountains of the Greater Caucasus with in Azerbaijan Republic. British Journal of Educational and Scientific Studies, 1. (23), January - June, 2016 VOLUME XII, Imperial College Press: 579-585. (in Eng.).

- [7] Guliyeva S.Y., Tarihazer S.A., Kuchinskaya I.Y. (2014) Geodynamic aspects of the formation of landslide landscapes in the north-eastern part of the Greater Caucasus (within Azerbaijan). In Materials of the geomorphologic congress. Volgograd: 80-86. (in Eng.).
- [8] John F., Shroder J.R., Brandon J. Weihs. (2010) Geomorphology of the lake Shewa landslide dam, Badakhshan, Afghanistan, using remote sensing data. *Geografiska Annaler: Series A, Physical Geography*, Volume 92, Issue 4, December: 469-483. (in Eng.).
- [9] Mardanov I.I. (2009) About possibilities of use of expert system for forecasting of landslides. In Works of Geographical society of Azerbaijan, volume XIV, Geographical problems of ensuring ecological safety of natural and economic systems, Baku: 36-40. (in Russian).
- [10] Mekhtiyev A.Sh. (1998) Space researches in Azerbaijan. In Materials of the second scientific and practical conference Azerbaijan on a Threshold of the XXI Century. Baku: 343-346. (in Russian)
- [11] Museibov M.A. (1998) Physical geography of Azerbaijan. The textbook for geographical faculties of universities. Baku, Maarif. (in Azeri).
- [12] Mardanov I.I., Hajizadeh F.M. (2003) The factors influencing landslide processes in Apsheron. Desertification problems in Azerbaijan // In Materials of the scientific and practical conference devoted to the 75 anniversary since the birth of the academician B. A. Budagov. Baku: 192-195. (in Russian).
- [13] Mardanov I.I., Aliyev E. A. (2016) Influence of the Geodynamic situation on ecotourism development in highlands of Azerbaijan part of Great Caucasus. *TOURISM ECONOMICS, The business and finance of tourism and recreation*, ISSN 1354-8166 (print); 2044-0375 (online). Vol. 22, 1.1, 2016, February: 94-100. (in Eng.).
- [14] Makarov M.I., Glaser B., Zech W., Malysheva T.I., Bulatnikova I.V., Volkov A.V. (2003) Nitrogen dynamics in alpine ecosystems of the northern Caucasus. *Plant and soil*, October, Vol. 256, Issue 2: 389. (in Eng.).
- [15] Mardanov I.I., Medzhidov D.B., Medzhidova V. G., Abdullaev G.G. (2005) The system analysis of the factors characterizing processes of an landslide forming in Apsheron. *ANASA News (It is devoted to day of astronautics) // Physics and technology problems of ways of remote sensing*. Vol. 8, 1 (8): 85-92. (in Russian).
- [16] Mehbaliev M.M., Jarullayev A.Sh., Mardanov I.I. (2016) Morphometric study and evaluation of erosion hazards of mountain geomorphosystem slopes // *Ciencia E Tecnica. Vitivinicola a science and technology journal*. ISSN: 0254-0223. Vol. 31 (2, 2016): 457-467. (in Eng.).
- [17] Pashayev N.A. (2007) Management of natural disasters in the territory of the Azerbaijan Republic. Works of Geographical society of Azerbaijan // *Modern geographical researches in Azerbaijan*. Vol. XI. Baku: 283-289. (in Russian).
- [18] Pike R.J. (2000) Geomorphometry diversity in quantitative surface analysis // *Progress in Physical Geography*, 24 (1): 1-20. (in Eng.).
- [19] Rainer Bell, Helene Petschko, Matthias Röhrs and Andreas Dix (2012) Assessment of landslide age, landslide persistence and human impact using airborne laser scanning digital terrain models // *Geografiska Annaler: Series A, Physical Geography Special Issue: Thematic Issue: Concepts and implications of environmental change and human impact: studies from Austrian geomorphological research*, March, Vol. 94, Issue 1: 135-156. (in Eng.).
- [20] Shary P.A., Sharaya L.S., Mitusov A.V. (2002) Fundamental quantitative methods of land surface analysis // *Geoderma*. Vol. 107, 1-2: 1-32. (in Eng.).
- [21] Shaw J., Sharpe D., Harris J. (2010) A flowline map of glaciated Canada based on remote sensing data // *Canadian Journal of Earth Sciences*, NRC Research Press, Vol. 47, 1: 89-101 (13). (in Eng.).
- [22] Taud H., Parrot J.F. (2005) Measurement of DEM roughness using the local fractal dimension // *Geomorphologie: relief, environnement*, 4: 327-338. (in Eng.).
- [23] Thompson J.A., Bell J.C., Butler C.A. (2001) Digital elevation model resolution: effects on terrain attribute calculation and quantitative soil-landscape modeling // *Geoderma*, 100: 67-89. (in Eng.).
- [24] Zhilin Li, Qing Zhu, Chris Gold (2005) Digital terrain modeling: principles and methodology. CRC PRESS. Boca Raton London New York Washington, D.C. 318 p. (in Eng.).
- [25] Zhou Q., Lees B., Tang G. (2008) Advances in Digital Terrain Analysis // *Lecture Notes in Geoinformation and Cartography Series*. Springer: 462 p. (in Eng.).

**И. И. Марданов¹, С. А. Тарихазер²,
И. Ю. Кучинская², С. Ю. Кулиева², Л. А. Исмаилова³**

¹СГУ, Сумгаит, Азербайджан,

²Институт Географии НАНА, Баку, Азербайджан,

³Азербайджанский государственный университет нефти и промышленности, Баку, Азербайджан

ГЕОСИСТЕМНЫЙ АНАЛИЗ РАЗВИТИЯ ОПОЛЗНЕВЫХ ЛАНДШАФТОВ ЮГО-ВОСТОЧНОГО СКЛОНА БОЛЬШОГО КАВКАЗА

Аннотация. Юго-восточная оконечность Большого Кавказа, входящая в пределы Азербайджана и являющаяся частью Альпийско-Гималайского геосинклинального пояса, характеризуется высокой динамичностью экзогенных рельефообразующих процессов, имеющих существенное влияние на развитие ландшафтной ситуации. В этой связи необходимо разработать различные методы для создания сценариев возможных изменений геоэкологической ситуации в различных массивах, имеющих пагубные последствия, путем выявления причинно-следственных связей. С этой целью в этой работе были проанализированы возможности прогнозирования оползней в природной области Большого Кавказа с использованием имеющихся геологических и геоморфологических, климатических и ландшафтных данных, а также визуальных наблюдений. Эти данные позволили выявить основные особенности факторов условий рельефа, гидроклиматические условия в некоторых физико-географических районах, в которых проявляются оползни, характер и интенсивность землепользования, в какой-то мере влияющие на сход оползней и их разрушительное воздействие.

Ключевые слова: процессы, склон, экзодинамический, ландшафты, эрозия, фактор, подрайон, Кавказ.

Information about authors:

Mardanov Ilham Ildirimoglu – Assistant of professor of branch “physical geography” Sumgait State University, Sumgait, Azerbaijan Republic, E-mail: geography.sumqayit@mail.ru,

Tarikhazer Stara Abulfas gyizi – candidate of geographical sciences, associate professor, leading research worker of Institute of Geography named by acad. H. A. Aliyev of ANAS, Azerbaijan, Baku, E-mail: kerimov17@gmail.com

Kuchinskaya Irina Yakovlevna – candidate of geographical sciences, associate professor, leading research worker of Institute of Geography named by acad. H. A. Aliyev of ANAS, Azerbaijan, Baku, E-mail: Irina.danula@gmail.com

Guliyeva Sevil Yunis gyizi – candidate of geographical sciences, associate professor, leading research worker of Institute of Geography named by acad. H. A. Aliyev of ANAS, E-mail: selinsevil@gmail.com

Ismaylova Latifa Arifgyizi – PhD student Azerbaijan State Oil and Industry University, Baku, Azerbaijan Baku, Azerbaijan, E-mail: Latifa.isamylova@gmail.com

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)

<http://www.bulletin-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 20.07.2018.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
16,25 п.л. Тираж 500. Заказ 4.