

ISSN 2518-1467 (Online),  
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Ш Ы С Ы

---

---

**ВЕСТНИК**

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

**THE BULLETIN**

THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

4

JULY – AUGUST 2019

---

---

ALMATY, NAS RK

---

*NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.*

*Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.*

*НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.*

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

**М. Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

**Абиев Р.Ш.** проф. (Ресей)  
**Абишев М.Е.** проф., корр.-мүшесі (Қазақстан)  
**Аврамов К.В.** проф. (Украина)  
**Аппель Юрген** проф. (Германия)  
**Баймуқанов Д.А.** проф., корр.-мүшесі (Қазақстан)  
**Байтулин И.О.** проф., академик (Қазақстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Қазақстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., РҒА академигі (Ресей)  
**Гашимзаде Ф.** проф., академик (Әзірбайжан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., корр.-мүшесі (Қазақстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Қалимолдаев М.Н.** проф., академик (Қазақстан), бас ред. орынбасары  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лупашку Ф.** проф., корр.-мүшесі (Молдова)  
**Мохд Хасан Селамат** проф. (Малайзия)  
**Мырхалықов Ж.У.** проф., академик (Қазақстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., корр.-мүшесі (Қазақстан)  
**Полещук О.Х.** проф. (Ресей)  
**Поняев А.И.** проф. (Ресей)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Қазақстан)  
**Таткеева Г.Г.** проф., корр.-мүшесі (Қазақстан)  
**Умбетаев И.** проф., академик (Қазақстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., РҒА корр.-мүшесі (Ресей)  
**Якубова М.М.** проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
<http://www.bulletin-science.kz/index.php/en/>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р  
д. х. н., проф. академик НАН РК  
**М. Ж. Журинов**

Р е д а к ц и о н н а я к о л л е г и я:

**Абиев Р.Ш.** проф. (Россия)  
**Абишев М.Е.** проф., член-корр. (Казахстан)  
**Аврамов К.В.** проф. (Украина)  
**Апель Юрген** проф. (Германия)  
**Баймуканов Д.А.** проф., чл.-корр. (Казахстан)  
**Байтулин И.О.** проф., академик (Казахстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Казахстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., академик РАН (Россия)  
**Гашимзаде Ф.** проф., академик (Азербайджан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., чл.-корр. (Казахстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Калимолдаев М.Н.** академик (Казахстан), зам. гл. ред.  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лунашку Ф.** проф., чл.-корр. (Молдова)  
**Моход Хасан Селамат** проф. (Малайзия)  
**Мырхалыков Ж.У.** проф., академик (Казахстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., чл.-корр. (Казахстан)  
**Полещук О.Х.** проф. (Россия)  
**Поняев А.И.** проф. (Россия)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Казахстан)  
**Таткеева Г.Г.** проф., чл.-корр. (Казахстан)  
**Умбетаев И.** проф., академик (Казахстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., член-корр. РАН (Россия)  
**Якубова М.М.** проф., академик (Таджикистан)

**«Вестник Национальной академии наук Республики Казахстан».**

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

**M. Zh. Zhurinov**

E d i t o r i a l b o a r d:

**Abiyev R.Sh.** prof. (Russia)  
**Abishev M.Ye.** prof., corr. member. (Kazakhstan)  
**Avramov K.V.** prof. (Ukraine)  
**Appel Jurgen,** prof. (Germany)  
**Baimukanov D.A.** prof., corr. member. (Kazakhstan)  
**Baitullin I.O.** prof., academician (Kazakhstan)  
**Joseph Banas,** prof. (Poland)  
**Bersimbayev R.I.** prof., academician (Kazakhstan)  
**Velesco S.,** prof. (Germany)  
**Velikhov Ye.P.** prof., academician of RAS (Russia)  
**Gashimzade F.** prof., academician ( Azerbaijan)  
**Goncharuk V.V.** prof., academician (Ukraine)  
**Davletov A.Ye.** prof., corr. member. (Kazakhstan)  
**Dzhrbashian R.T.** prof., academician (Armenia)  
**Kalimoldayev M.N.** prof., academician (Kazakhstan), deputy editor in chief  
**Laverov N.P.** prof., academician of RAS (Russia)  
**Lupashku F.** prof., corr. member. (Moldova)  
**Mohd Hassan Selamat,** prof. (Malaysia)  
**Myrkhalykov Zh.U.** prof., academician (Kazakhstan)  
**Nowak Isabella,** prof. (Poland)  
**Ogar N.P.** prof., corr. member. (Kazakhstan)  
**Poleshchuk O.Kh.** prof. (Russia)  
**Ponyaev A.I.** prof. (Russia)  
**Sagiyani A.S.** prof., academician (Armenia)  
**Satubaldin S.S.** prof., academician (Kazakhstan)  
**Tatkeyeva G.G.** prof., corr. member. (Kazakhstan)  
**Umbetayev I.** prof., academician (Kazakhstan)  
**Khripunov G.S.** prof. (Ukraine)  
**Yuldashbayev Y.A.,** prof. corresponding member of RAS (Russia)  
**Yakubova M.M.** prof., academician (Tadjikistan)

**Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.**

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

UDS 581.1:34.05.15

**K. R. Uteulin<sup>1</sup>, O. A. Agapov<sup>2</sup>, K. Z. Zambakin<sup>1</sup>, Y. T. Martirosyan<sup>3</sup>**<sup>1</sup>Republican state enterprise «Institute of Plant Biology and Biotechnology, Committee of Sciences,  
Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan,<sup>2</sup>Republican state enterprise “Scientific-Research Center “Garysh-Ecology” Aerospace committee,  
the Ministry of Defense and Aerospace Industry of the Republic of Kazakhstan, Almaty, Kazakhstan,<sup>3</sup>Federal State Budgetary Institution All-Russian Research Institute of Agricultural Biotechnology, Moscow, Russia.

E-mail: gen\_uteulink@mail.ru, agaole@mail.ru, zhambakin@gmail.com, yumart@yandex.ru

## OPTIMAL MOISTURE OF SOIL SAMPLES FOR SEED GERMINATION AND GROWTH OF SEEDLINGS IN CLOSED CONTAINERS

**Abstract.** The purpose of the research is to determine optimum moisture content of soil samples in closed, transparent containers for seed germination and seedling growth of wild-growing wormwood – artemisia effusus (*Artemisia diffusa*), wild-growing type of cereal – comb-shaped wheat grass (*Agropyron pectiniforme*), cultured cereal – soft wheat (*Triticum aestivum*), variety Saratovskaya-29. Tested soil samples with humidity of 20%, 40% and 60%.

It is established that in a closed container (5 liters) soil moisture in the range of 40% - 60% provides the best seed germination and the growth of seedlings of the used plant species.

In tested closed containers, soil moisture of 40% should be used, with relatively short period of plants research (30 days). With a longer period of research, it is possible to use soil with moisture content of up to 60%, because with time, the excess water is absorbed by growing plants.

The method of closed containers find use for studies of influence of toxic, volatile, gaseous substances on the growth and development of plants in laboratory conditions. The results of this work will be used in modeling the ecosystem of Central Kazakhstan in connection with environmental pollution of volatile, highly toxic components of rocket-engine propellant, resulting from activities of Baikonur space-launch complex.

**Key words:** closed containers, soil moisture, seed germination, growth of seedlings.

**Introduction.** The issues of influence volatile, toxic substances on the growth and development of plants and accumulation of volatile toxic substances by wild and cultivated plant species are of high concern. For example, plants of Central Kazakhstan are under the polluting influence of unspent rocket-engine propellant - unsymmetrical dimethyl hydrazine (UDMH) and its transformation products. UDMH and its transformation products enters the plants and then into the body of animals and humans from soil and through aerogenic way [1]. Volatile derivatives of UDH have been identified in various types of soil [2]. As is known, UDH and a number of its derivatives belong to the first class of hazard (extremely dangerous organic compound) [3].

Volatile toxic substances are chemicals that could be available in the air in a gaseous state. In this regard, to estimate phytotoxicity of UDMH and its derivatives under laboratory conditions should be used methods, similar to those for study of gas resistance of the plants.

It is known that (closed containers, ‘gas chambers’) are used to study gas resistance of the plants in laboratory conditions. Plant or its parts are placed in container, gases are dispensed (for example, sulfur dioxide, hydrogen fluoride and hydrogen chloride) and then changes in plant reactions are monitored. As diagnostic features utilized the following parameters: change in germination and seed germination energy, growth rate and formation of individual organs, change in the timing and duration of the developmental and organogenesis phases, and others. In the event of activity of sublethal and lethal gas concentrations, plant resistance can be determined by degree of leaf damage necrosis or by changes in the water regime and photosynthesis [4-7].

One of the first task in development of technology for evaluating the effect of volatile toxic substances on plants in closed containers is to establish optimum soil moisture, which provides the best seed germination and development of plant seedlings.

**Purpose of the article** – establish the optimum moisture content of soil samples in closed container for seed germination and growth of seedlings of wormwood and cereals.

**Methods.** The targets of research were wild-growing wormwood: artemisia effusus (*Artemisia diffusa*), wild-growing type of cereal - comb-shaped wheat grass (*Agropyron pectiniforme*), cultivated type of cereal - soft wheat (*Triticum aestivum*), variety Saratovskaya-29.

In work process being used ready-made soil universal, for all types of vegetable, green, flower crops and seedlings, environmentally friendly product 'Himself agronomist'. Production Russia, LLC 'AgroSnabRetail'. Composition: drug turf (lime carbonate, chalk, dolomite powder) and structuring materials, mineral fertilizers.

Content of nutrients, mg/ml:	Nitrogen (NH <sub>4</sub> +NO <sub>3</sub> )	240-350
	Phosphorus (P <sub>2</sub> O <sub>5</sub> )	290-410
	Potassium (K <sub>2</sub> O)	330-470
	Acidity pH	5,4-6,6

Complete set of required of minor elements.

Ready-made universal soil was dried at room temperature for two weeks; sand was thoroughly washed and dried. Prepared a mixture of soil and sand with weight ratio of 70% soil and 30% sand. Sand was added to better aerate the soil. Then prepared soil samples with different humidity: 20%, 40% and 60%.

Transparent plastic bottles (commercial, household, used for drinking water) of 5.25 liters were cut horizontally centrally, 0.5 kg prepared soil samples placed in containers. The seeds were sown, then containers sealed with scotch tape and tightly closed with cap on top (figure).



Seed germination and seedling growth of comb-shaped wheat grass (*Agropyron pectiniforme*) in closed containers

Seed germination was carried out at temperature of + 20 + 23 °C.

Used known methods of seed germination [8]. Each batch of each series was carried out in three replications, followed by statistical processing [9].

**Results and discussion.** Being studied seed germination of wild-growing wormwood – artemisia effusus (*Artemisia diffusa*) in closed containers with soil samples of different moisture. The seeds of wormwood are small (2x0.8 mm), lightweight (weight of 1,000 seeds is 0.2-0.3 grams). 0.5 grams of wormwood seeds were sown, or about 1,500 seeds in each container. The observation period was 30 days. After 30 days, containers were opened, the best condition of seedlings was found at soil sample with 40% of moisture, soil sample with 60% moisture was waterlogged.

Seed germination of wormwood was 1%, 1.6% and 1.7% on soil samples with moisture content of 20%, 40% and 60% respectively. Seed germination of wormwood is 1,6-1,7 times higher on soil samples of 40% and 60% compared to soil samples with moisture content of 20% (table 1).

Table 1 – Seed germination of artemisia effusus (*Artemisia diffusa*) in closed containers within one month. 0.5 grams of seeds or about 1,500 seeds were sown in each container. Represented data collected on 30th day of research

Soil moisture, %	Germination, %	Seedling height, cm
20	1,0±0,02	1,2 ±0,3
40	1,6±0,03	3,2±0,9
60	1,7±0,03	3,1±1,0

Seeds of wormwood were collected in November 2018, the period of post-harvest ripening was 3,5 months. Such a short period of post-harvest ripening explains the low germination of wormwood seeds.

Part of containers were opened later, only after 3 months. It has been established that condition of container with 60% soil moisture is significantly improved. As far as water is used to increase biomass of growing plants and its content in soil and air is reduced.

The results of research cereal seeds are presented in tables 2 and 3, respectively.

Table 2 – Seed germination and seedling growth of comb-shaped wheat grass (*Agropyron pectiniforme*) in closed containers. 300 seeds are sown in each container. Represented data collected on 30th day of research

Soil moisture, %	Germination, %	Seedling height, cm
20	3,6±0,03	15 ±3
40	42±1,2	17±4
60	41±1,3	18±4

Table 3 – Growth and development of soft wheat seedlings (*Triticum aestivum*), variety Saratovskaya-29 in closed containers within one month. In each container, 100 seeds are sown. Represented data collected on 30th day of research

Soil moisture, %	Germination, %	Seedling height, cm
20	19±0,8	27 ±4
40	78±2,2	37±5
60	79±2,3	37±6

Research of cereal seeds germination was carried out during a month; it was established that soil moisture of 40% is optimal for seed germination and seedling growth in closed containers. Soil samples with moisture content of 20% had significantly lower seed germination compared to seed germination of soil samples with moisture content of 40% and 60%. The germination of cereal seeds and seedling development was about the same of soil samples with moisture content of 40% and 60%. (tables 2 and 3).

In closed containers, both for wormwood and for cereals, soil moisture in range of 40-60% provides the best seed germination. Seeds of wild plant species are heterogenic and do not germinate simultaneously. Then, cultivated cereal, soft wheat (*Triticum aestivum*) variety Saratovskaya-29 is able to complete seed germination on 14th day of research. Quite opposite seeds of wild-growing species of comb-shaped wheat grass (*Agropyron pectiniforme*) and artemisia effusus (*Artemisia diffusa*) begin to germinate on 3-4 days or later on the 14th day.

Regarding the optimal moisture content of soil samples in closed containers, for comparison, we quote data of Pryanishnikov D. N. (1900) about optimum moisture content of open ground (soil) for seed germination, plant growth and development. In open ground, the lower limit of optimum moisture content is approximately estimated depending on the type of soil and plants by the following values: for grasses - 50–60%, for cereals - 45–50%, for vegetable and technical crops - 40–45% [10].

That is, there is an approximate coincidence of the optimum soil moisture for plant development both in open ground and in closed container.

However, the results of our research have shown waterlogged general condition of closed container with soil sample of 60% humidity. Waterlogging often leads to mold and contamination comparing with soil samples having less moisture. It is known that the development of bacteria stops at about 25% of environment moisture and about 15% for mold [11].



**Conclusions.** In closed, transparent container (5 liters) soil moisture (soil mixed with peat and sand) in the range of 40–60% compared to humidity of 20% provides the best seed germination of wild species: wormwood (*Artemisia diffusa*), comb-shaped wheat grass (*Agropyron pectiniforme*), cultivated cereal soft wheat (*Triticum aestivum*) variety Saratovskaya-29.

In tested closed containers, soil moisture of 40% should be used, with relatively short research period of plants (30 days). With a longer period of research, it is possible to use soil with moisture content of up to 60%, because with time, the excess water is absorbed by growing plants.

The method of closed containers find use for studies of influence of toxic, volatile, gaseous substances on the growth and development of plants in laboratory conditions. The results of this work will be used in modeling the ecosystem of Central Kazakhstan in connection with environmental pollution of volatile, highly toxic components of rocket-engine propellant, resulting from activities of Baikonur space-launch complex.

**Funding. Source of research funding.** Republican budget program 008 'Applied research in the field of space activities'. Scientific and technical project: 'Study of the nature of plant pollution by unsymmetrical dimethyl hydrazine and toxic products of its transformation'. Republican state enterprise. Research Center "Garysh-Ecology". Aerospace Committee of the Ministry of Defense and Aerospace Industry of the Republic of Kazakhstan.

К. Р. Утеулин<sup>1</sup>, О. А. Агапов<sup>2</sup>, К. Ж. Жамбакин<sup>1</sup>, Ю. Ц. Мартиросян<sup>3</sup>

<sup>1</sup>«Биология және өсімдіктер биотехнологиясы институты» Республикалық мемлекеттік кәсіпорны, Қазақстан Республикасының Білім және ғылым министрлігі, Алматы, Қазақстан,

<sup>2</sup>Қазақстан Республикасы Қорғаныс және аэроғарыш өнеркәсібі министрлігі Аэроғарыш комитеті, «Ғарыш-Экология» ғылыми-зерттеу орталығы» Республикалық мемлекеттік кәсіпорны, Алматы, Қазақстан,

<sup>3</sup>Ауылшаруашылық биотехнологиясының Бүкілресейлік ғылыми-зерттеу институты, Мәскеу, Ресей

#### ТҰҚЫМДАРДЫҢ ЖӘНЕ ТҰҚЫМ ӨСКІНДЕРІНІҢ ЖАБЫҚ ЫДЫСТАРДА ӨСУІ ҮШІН ТОПЫРАҚТАР ҮЛГІСІНІҢ ОҢТАЙЛЫ ЫЛҒАЛДЫЛЫҒЫ

**Аннотация.** Зерттеу мақсаты жусанның жабайы өсетін түрінің тұқымдарының және екпе көшеттердің өсуі үшін топырақ үлгісінің жабық, мөлдір ыдыстардағы оңтайлы ылғалдылығын белгілеу болды - тармақты жусан (*Artemisia diffusa*), дақылдың жабайы өсетін түрі - еркек бидайық (*Agropyron pectiniforme*), мәдениленген дақыл - жұмсақ бидай (*Triticum aestivum*), Саратов сұрыбы-29. 20, 40 және 60 % ылғалдылықтағы топырақ үлгілері сыналды. Жабық ыдыста (5 литрлік) топырақтың ылғалдылығы 40-60 % интервалда тұқымдардың және екпе көшеттердің пайдаланылған өсімдік түрлерінде жақсы өсуін қамтамасыз ететіні анықталған. Сыналған жабық ыдыстарда өсімдіктердің қатыстық қысқа кезеңінде 40 % тең топырақ үлгісінің ылғалдылығын пайдалану қажет (30 тәулік). Пайдаланудың неғұрлым ұзақ кезеңінде 60 % дейінгі ылғалдылықпен топырақты пайдалану мүмкіндігі бар, себебі уақыт өте келе судың артығы өсіп келе жатқан өсімдікпен игеріледі. Жабық ыдыстар әдісі зертханалық жағдайда өсімдіктердің өсуіне және дамуына улы, ұшатын, газ тәрізді заттардың әсер етуін зертеуде қолданылады. Осы жұмыстардың нәтижелері Орталық Қазақстанның экожүйесін Байқоңыр космодромының қызметіне байланысты қоршаған ортаның ұшқыш, жоғары токсинді ракеталық жанармаймен ластануына байланысты үлгілеуде пайдаланылатын болады.

**Түйін сөздер:** жабық ыдыстар, топырақтың ылғалдылығы, тұқымдардың өсуі, екпе көшеттердің өскіні.

К. Р. Утеулин<sup>1</sup>, О. А. Агапов<sup>2</sup>, К. Ж. Жамбакин<sup>1</sup>, Ю. Ц. Мартиросян<sup>3</sup>

<sup>1</sup>РГП «Институт биологии и биотехнологии растений», Алматы, Казахстан,

<sup>2</sup>РГП «Научно-исследовательский центр «Ғарыш-Экология», Алматы, Казахстан,

<sup>3</sup>Федеральное государственное бюджетное научное учреждение,

Всероссийский научно-исследовательский институт сельскохозяйственной биотехнологии, Москва, Россия

#### ОПТИМАЛЬНАЯ ВЛАЖНОСТЬ ПОЧВЫ ДЛЯ ПРОРАСТАНИЯ СЕМЯН И РОСТА СЕЯНЦЕВ В ЗАМКНУТЫХ ЕМКОСТЯХ

**Аннотация.** Целью исследований служило установление в замкнутых, прозрачных емкостях оптимальной влажности образцов почвы для всхожести семян и роста сеянцев дикорастущего вида полыни – полынь развесистая (*Artemisia diffusa*), дикорастущего вида злака - пырея гребневидного (*Agropyron pectiniforme*), окультуренного злака - мягкая пшеница (*Triticum aestivum*), сорт Саратовская-29. Использован универсальный, коммерческий торфогрунт.

Установлено, что в замкнутой, прозрачной емкости (на 5 литров) влажность почвы в интервале 40-60 % в сравнении с влажностью почвы 20 %, обеспечивает лучшую всхожесть семян и рост сеянцев *Artemisia diffusa*, *Agropyron pectiniforme* и *Triticum aestivum*, сорт Саратовская-29.

В испытанных замкнутых емкостях следует использовать влажность образцов почвы, равную 40 %, при относительно коротком периоде исследований растений (30 суток). При более длительном периоде исследований возможно использование почвы с влажностью до 60 %, по сколько со временем излишек воды усваивается растущими растениями.

Замкнутая, прозрачная емкость может служить моделью замкнутой экосистемы, то есть системы, не предполагающей какого-либо обмена веществом с внешней средой замкнутых емкостей. Такие системы представляют научный интерес и могут быть использованы для исследования влияния летучих (газообразных) соединений на растения.

Область применения: метод замкнутых емкостей находит применение в исследовании влияния летучих, газообразных веществ на рост и развитие растений в лабораторных условиях. Результаты этой работы будут использованы при моделировании экосистемы Центрального Казахстана в связи с загрязнением окружающей среды летучими, высокотоксичными компонентами ракетного топлива, возникающими в результате деятельности космодрома Байконур.

**Ключевые слова:** замкнутые емкости, проростание семян, влажность почвы.

#### Information about authors:

Uteulin Kairat, Head of the department. Associate professor, Institute of Plant Biology and Biotechnology, Committee of Sciences, Ministry of Education and Science of the Republic of Kazakhstan, Almaty, Kazakhstan; gen\_uteulink@mail.ru; <https://orcid.org/0000-0002-5459-0902>

Agapov Oles, Acting Head of the Department of Environmental Programs, RSE «Scientific-research center «Garysh-Ecology» (RSE «SRC «Garysh-Ecology»), Almaty, Kazakhstan; agaole@mail.ru; <https://orcid.org/0000-0003-3395-1358>

Zhambakin Kabyl, Professor, Member of NAS RK, Director General, Institute of Plant Biology and Biotechnology, Almaty, Kazakhstan; zhambakin@gmail.com; <https://orcid.org/0000-0001-5243-145X>

Martirosyan Y.T., Head of the group of aeroponic plant growing technologies, Federal State Budgetary Institution All-Russian Research Institute of Agricultural Biotechnology, Moscow, Russia; yumart@yandex.ru; <https://orcid.org/0000-0001-8825-238>

#### REFERENCES

- [1] Agapov O.A., Fedorina O.A., Atigaev A.B., Uteulin K.R., Zheksenbay A., Kazkeev D., Aldasugurova Ch.Zh., Kurbatova N.B. (2019) The effect of unsymmetrical dimethyl hydrazine on seed germination, growth and anatomical parameters of seedlings of wild plant species // Science News of Kazakhstan. N 1(139). P. 210-222 (in Rus.).
- [2] Kenesov B.N. (2008) Identification of volatile transformation products of unsymmetrical dimethyl hydrazine in soils using vapor phase extraction method in combination with chromatography-mass spectrometry // News of the National academy of sciences of the Republic of Kazakhstan. Chemical series. N 5. P. 48-53 (in Rus.).
- [3] GOST 12.1.005-76 (1989) Occupational safety standards system. Air working area. General sanitary and hygienic requirements. Introduction date. Developed and entered by the USSR Ministry of Health, All-Union Central Council of Trade Unions. 01.01.1989 (in Rus.).
- [4] Yudin Yu.N., Bezmenov A.Ya., Klyukina L.K., Mironov O.D. (1982) Method of evaluation gas resistance of plants. SU 1266490 A1 (in Rus.).
- [5] Yermeeva V.G., Denisova E.S. (2011) Gas resistance of woody plants of Siberia // Siberian Journal of Ecology. 2. P. 263-271 (in Rus.).
- [6] Fedorova A.I., Nikolskaya A.N. (2003) Workshop on ecology and environmental protection. M.: Vlados. 286 p. (in Rus.).
- [7] Fedulov Yu.P. (2015) Methods for determining plant resistance: a course of lectures. Krasnodar: Kuban state agrarian university. 39 p. (in Rus.).
- [8] Karpin V.I., Perepravo N.I., Zolotaryev V.N., Ryabova B.E., Shamsutdinova E.Z., Kozlova T.B. (2012) Method of determining the growth force of feed crops seeds. M.: Printing house Moscow Agricultural Academy named after K. A. Timiryazev. 16 p. (in Rus.).
- [9] Lakin G.F. (1990) Biometrics. M.: Printing house "Higher education". 352 p. (in Rus.).
- [10] Pryanishnikov D.N. (1990) About the effect of soil moisture on plant development // Journal of Experimental Agronomy. SPb. 1. P. 1-20 (in Rus.).
- [11] Orlov V.I. (1964) Fundamentals of microbiology and food hygiene: Textbook. M.: Economy. 208 p. (in Rus.).

---

---

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www.nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*  
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 19.07.2019.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
15,5 п.л. Тираж 500. Заказ 4.