

ISSN 2518-1467 (Online),  
ISSN 1991-3494 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Ш Ы С Ы

---

---

**ВЕСТНИК**

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

**THE BULLETIN**

THE NATIONAL ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

PUBLISHED SINCE 1944

5

SEPTEMBER – OCTOBER 2019

---

---

ALMATY, NAS RK

---

---

*NAS RK is pleased to announce that Bulletin of NAS RK scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of Bulletin of NAS RK in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential multidiscipline content to our community.*

*Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабаршысы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабаршысының Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді мультидисциплинарлы контентке адалдығымызды білдіреді.*

*НАН РК сообщает, что научный журнал «Вестник НАН РК» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Вестника НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному мультидисциплинарному контенту для нашего сообщества.*

Б а с р е д а к т о р ы

х. ғ. д., проф., ҚР ҰҒА академигі

**М. Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

**Абиев Р.Ш.** проф. (Ресей)  
**Абишев М.Е.** проф., корр.-мүшесі (Қазақстан)  
**Аврамов К.В.** проф. (Украина)  
**Аппель Юрген** проф. (Германия)  
**Баймуқанов Д.А.** проф., корр.-мүшесі (Қазақстан)  
**Байтулин И.О.** проф., академик (Қазақстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Қазақстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., РҒА академигі (Ресей)  
**Гашимзаде Ф.** проф., академик (Әзірбайжан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., корр.-мүшесі (Қазақстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Қалимолдаев М.Н.** проф., академик (Қазақстан), бас ред. орынбасары  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лупашку Ф.** проф., корр.-мүшесі (Молдова)  
**Мохд Хасан Селамат** проф. (Малайзия)  
**Мырхалықов Ж.У.** проф., академик (Қазақстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., корр.-мүшесі (Қазақстан)  
**Полещук О.Х.** проф. (Ресей)  
**Поняев А.И.** проф. (Ресей)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Қазақстан)  
**Таткеева Г.Г.** проф., корр.-мүшесі (Қазақстан)  
**Умбетаев И.** проф., академик (Қазақстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., РҒА корр.-мүшесі (Ресей)  
**Якубова М.М.** проф., академик (Тәжікстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабаршысы».

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы»РҚБ (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде  
01.06.2006 ж. берілген №5551-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.

Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
<http://www.bulletin-science.kz/index.php/en/>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор  
д. х. н., проф. академик НАН РК  
**М. Ж. Журинов**

Редакционная коллегия:

**Абиев Р.Ш.** проф. (Россия)  
**Абишев М.Е.** проф., член-корр. (Казахстан)  
**Аврамов К.В.** проф. (Украина)  
**Апель Юрген** проф. (Германия)  
**Баймуканов Д.А.** проф., чл.-корр. (Казахстан)  
**Байтулин И.О.** проф., академик (Казахстан)  
**Банас Иозеф** проф. (Польша)  
**Берсимбаев Р.И.** проф., академик (Казахстан)  
**Велесько С.** проф. (Германия)  
**Велихов Е.П.** проф., академик РАН (Россия)  
**Гашимзаде Ф.** проф., академик (Азербайджан)  
**Гончарук В.В.** проф., академик (Украина)  
**Давлетов А.Е.** проф., чл.-корр. (Казахстан)  
**Джрбашян Р.Т.** проф., академик (Армения)  
**Калимолдаев М.Н.** академик (Казахстан), зам. гл. ред.  
**Лаверов Н.П.** проф., академик РАН (Россия)  
**Лунашку Ф.** проф., чл.-корр. (Молдова)  
**Моход Хасан Селамат** проф. (Малайзия)  
**Мырхалыков Ж.У.** проф., академик (Казахстан)  
**Новак Изабелла** проф. (Польша)  
**Огарь Н.П.** проф., чл.-корр. (Казахстан)  
**Полещук О.Х.** проф. (Россия)  
**Поняев А.И.** проф. (Россия)  
**Сагиян А.С.** проф., академик (Армения)  
**Сатубалдин С.С.** проф., академик (Казахстан)  
**Таткеева Г.Г.** проф., чл.-корр. (Казахстан)  
**Умбетаев И.** проф., академик (Казахстан)  
**Хрипунов Г.С.** проф. (Украина)  
**Юлдашбаев Ю.А.** проф., член-корр. РАН (Россия)  
**Якубова М.М.** проф., академик (Таджикистан)

«Вестник Национальной академии наук Республики Казахстан».

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5551-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год

Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18.

www: nauka-nanrk.kz, bulletin-science.kz

---

© Национальная академия наук Республики Казахстан, 2019

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of chemistry, professor, academician of NAS RK

**M. Zh. Zhurinov**

E d i t o r i a l b o a r d:

**Abiyev R.Sh.** prof. (Russia)  
**Abishev M.Ye.** prof., corr. member. (Kazakhstan)  
**Avramov K.V.** prof. (Ukraine)  
**Appel Jurgen,** prof. (Germany)  
**Baimukanov D.A.** prof., corr. member. (Kazakhstan)  
**Baitullin I.O.** prof., academician (Kazakhstan)  
**Joseph Banas,** prof. (Poland)  
**Bersimbayev R.I.** prof., academician (Kazakhstan)  
**Velesco S.,** prof. (Germany)  
**Velikhov Ye.P.** prof., academician of RAS (Russia)  
**Gashimzade F.** prof., academician ( Azerbaijan)  
**Goncharuk V.V.** prof., academician (Ukraine)  
**Davletov A.Ye.** prof., corr. member. (Kazakhstan)  
**Dzhrbashian R.T.** prof., academician (Armenia)  
**Kalimoldayev M.N.** prof., academician (Kazakhstan), deputy editor in chief  
**Laverov N.P.** prof., academician of RAS (Russia)  
**Lupashku F.** prof., corr. member. (Moldova)  
**Mohd Hassan Selamat,** prof. (Malaysia)  
**Myrkhalykov Zh.U.** prof., academician (Kazakhstan)  
**Nowak Isabella,** prof. (Poland)  
**Ogar N.P.** prof., corr. member. (Kazakhstan)  
**Poleshchuk O.Kh.** prof. (Russia)  
**Ponyaev A.I.** prof. (Russia)  
**Sagiyani A.S.** prof., academician (Armenia)  
**Satubaldin S.S.** prof., academician (Kazakhstan)  
**Tatkeyeva G.G.** prof., corr. member. (Kazakhstan)  
**Umbetayev I.** prof., academician (Kazakhstan)  
**Khripunov G.S.** prof. (Ukraine)  
**Yuldashbayev Y.A.,** prof. corresponding member of RAS (Russia)  
**Yakubova M.M.** prof., academician (Tadjikistan)

**Bulletin of the National Academy of Sciences of the Republic of Kazakhstan.**

**ISSN 2518-1467 (Online),**

**ISSN 1991-3494 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5551-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/>, <http://bulletin-science.kz>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

S. A. Grikshas<sup>1</sup>, V. V. Kalashnikov<sup>2</sup>, G. K. Dzhanibekova<sup>3</sup>,  
G. A. Funikov<sup>1</sup>, A. V. Ovchinnikov<sup>1</sup>, N. I. Kulmakova<sup>1</sup>, V. T. Yepifanov<sup>1</sup>,  
V. V. Khramtsov<sup>1</sup>, S. N. Sarimbekova<sup>3</sup>, M. Sh. Yerezhepova<sup>3</sup>

<sup>1</sup>FSBEI HE Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev,  
Moscow, Russia,

<sup>2</sup>Federal state budgetary institution “All-Russian Scientific Research Institute of Horse Breeding”,  
Divovo village, Rybnovsky District, Ryazan Region, Russia,

<sup>3</sup>Kazakh National Agrarian University, Almaty, Republic of Kazakhstan.

E-mail: stepangr56@mail.ru, vniik08@mail.ru, guka.67@mail.ru, gafvt@yandex.ru,  
svinovod@rgau-msha.ru, knil1@mail.ru, manager@nauchnik.org,  
komarovaoksana572@gmail.com, sarimbekova@yandex.ru, princessa.maki@mail.ru

## PRODUCTIVITY AND BIOLOGICAL FEATURES OF PIGS OF DOMESTIC AND CANADIAN BREEDING

**Abstract.** In the conditions of large pig breeding complexes in the raw material base of Smolmyaso OJSC and Pavlovskaya Sloboda Meat Processing Plant OJSC, in the period of 2010-2015, research and production experience was conducted to study the productivity and biological features of domestic and western breeding pigs. Experimental young stock from 2 months age till reaching the live weight of 100 kg was kept in the control growing. The conditions of keeping, feeding and slaughter of young pigs were identical. When young pigs reached 95-105 kg in live weight, they were evaluated by their own productivity and slaughter indices.

In the process of research, it was found that among the domestic and Canadian young stock, the highest precocity and growth rate was characterized by three-breed hybrid young stock, obtained by crossing hybrid sows with Duroc boars. The results of the control slaughter showed that among the domestic and Canadian young stock, the best slaughter indices were characterized by the three-breed hybrid young.

It was established that the three-breed cross young pigs of the Canadian breeding reached the pre-slaughter live weight 22 days earlier compared to the domestic one, the average daily weight gain was 134.1 g higher, the slaughter yield was 2.3% higher. Thus, the three-breed commercial young stock of Canadian breeding reliably surpasses the pigs of domestic breeding.

The calculation of the economic efficiency of growing young pigs shows that the additional income from the sale of 1 head of Canadian breeding is 1.5 thousand rubles (23 US dollars) compared with the domestic breeding.

Based on the research results, it was recommended that pig farms engaged in breeding and growing domestic pigs, should strengthen breeding to improve the fattening and meat qualities of pigs. To obtain commercial young pigs with high fattening and meat qualities, it is recommended to use Canadian pigs more extensively.

**Keywords:** young pigs, domestic breeding, Canadian breeding, growth rate, average daily gains, slaughter weight, slaughter yield.

**Introduction.** Currently, animal husbandry is one of the most important sectors of the agro-industrial complex of the Russian Federation and plays a significant role in the country's food supply. Pig breeding, as the most intensive and efficient livestock industry, makes a substantial contribution to providing the population of the country with meat and meat products. At the moment, the pig breeding is developing dynamically, which means that the production of pork sustainably develops. As before, the growth rate of pork production goes ahead of the increase in livestock population, which indicates the intensification of the industry due to the implementation of progressive methods of pig breeding, the involvement of highly productive breeds in the production and widespread use of hybridization and crossing, and the improvement of fattening technology and pig breeding [1-3].

According to the National Union of pig breeders, for the last 5 years from 2014 to 2018, pork production increased from 2816 thous. t to 3733 thous. t. or by 24.6%. The main growth in pork production was received due to the increase in the production of agricultural enterprises. During this period, the import of pork dropped from 427 thous. t to 80 thous. t or 5.3 times [4]. The pork production figures show that the pig breeding successfully executes the program of import substitution.

A significant portion of pork is produced in the large industrial complexes with the use of intensive technologies of growing and fattening animals. However, as indicated by Yu.V. Tatulov and co-authors [5, 6], instability of dimensional and weight standards of pigs, no stress resistance lead to a decrease in quality and appearance of meat defects, complicating its industrial processing. In this regard, there is an ongoing search for breeds, lines and types of pigs that are able to show high productive quality in large pig-breeding complexes [7-10].

Currently, the domestic pig farms to form tribal flocks often use imported pigs from abroad and, in particular, from Canada. These pigs are characterized by high productive qualities [12-17].

But so far the lifetime productivity and biological features of Canadian breeding pigs in large farms are poorly studied.

Therefore, the conduct of a comparative assessment of productivity and biological features of domestic and Canadian pigs in the conditions of large pig farms is very urgent.

**The aim of the research** is a determination of biological and productive features in pigs of domestic and the Canadian breeding grown in the conditions of the pig farms.

To achieve this aim, the following objectives were traced:

- to assess the own productivity of young pigs of domestic breeding;
- to assess the own productivity of young pigs of the Canadian breeding;
- to determine the slaughter parameters of the tested young stock;
- to calculate the economic efficiency of pork production when using domestic and Canadian pigs.

**Arrangement and methods of the research.** The control growing of experimental young pigs and slaughter were carried out in the raw material zone of Smolmya so OJSC and Pavlovskaya Sloboda Meat Processing Plant OJSC of the Moscow Region during 2010-2015.

To conduct a research and production experiment, two experimental groups were formed with 3 subgroups of the most common pig breeds and breed combinations. The first group consists of yelts from parents of domestic breeding, in the second - from Canada breeding. The sequence of the experiment is shown in scheme.

In all experiments, identical methods of housing and feeding, as well as transportation and pre-slaughter preparation and slaughter of animals were applied. Pigs were fed according to the norms and diets of feeding at the animal farms [18].

<b>Formation of the experimental groups</b>	
Domestic breeding: 1.1. Large white x Large white 1.2. Large white x Landrace 1.3. (Large white x Landrace) x Duroc	Canadian breeding: 2.1. Yorkshire 2.2. Yorkshire x Landrace 2.3. (Yorkshire x Landrace) x Duroc
Test indicators	
At control growing	At slaughter
Pre-slaughter live weight, kg Age of reaching 100 kg mass, days Average daily weight gain, g	Carcass weight, kg Mass of internal fat, kg Deadweight, kg Slaughter yield, %
Calculation of economic efficiency.	

Scheme of the experiment performance

Lifetime productivity of experimental young stock was assessed according to standard methods for control growing according to indicators covering the animals' productivity - age of attaining live weight in days and average daily gain in grams [19].

Slaughter indicators were determined in accordance with the "Methodological recommendations by the V.I. Lenin Academy of Agricultural Sciences on the assessment of meat productivity, meat quality and subcutaneous fat of pigs" [19].

Evaluation of the economic efficiency of growing experimental young pigs was determined by mapping live weight during realization and the costs of growing young pigs in monetary terms. The level of profitability was calculated as the ratio of profit to prime cost, %.

Biometric processing of the obtained data was performed according to A.M. Gataulin's methodological guidelines on the design of measurement results using the Microsoft Excel, the accuracy of the difference was taken at the reliability threshold of  $B1 = 0.95$  (significance level  $P \leq 0.05$ ). At the level of difference  $P \geq 0.05$ , the difference is not statistically significant [20]. As a control group, young pigs of 1.1 and 2.1 subgroups were used.

**Research results and discussion.** Lifetime assessment of young pigs in their own productivity allows to evaluate animals according to phenotype. The results of table 1 show that among domestic breeding pigs, pre-slaughter mass ranged from 99.3 kg (subgroup 1.1) to 102.3 kg (subgroup 1.3), which indicates slight deflections in this indicator.

In the three-breed cross young stock of domestic breeding from 1.3 subgroup, the age of reaching 100 kg of live weight was 178 days, which is less compared to gilts from 1.1 and 1.2 subgroups by 8 ( $P \leq 0.05$ ) and 4 days respectively. The highest gains in live weight were obtained from gilts of 1.3 subgroup - 694.0 g, which is higher than in 1.1 and 1.2 subgroups by 52.0 g or 7.5% ( $P \leq 0.01$ ) and 21.9 g - 3.2%, respectively.

Consequently, among young stock of domestic breeding, the highest early ripeness and growth rate were for the three-crossbred stock, received by crossing crossbred sows with duroc boars.

Analysis of the data in Table 1 demonstrates that among young Canadian breeding pigs, the smallest age to reach 100 kg of weight was obtained from gilts of 2.3 group — 156 days, which is less compared to the young pigs from 2.1 and 2.2 subgroups by 6 ( $P \leq 0.05$ ) and 2 days respectively. The highest average daily gains of live weight among young Canadian breeding were obtained from gilts of 2.3 subgroup - 828.1 g, which is higher than in yelts from 2.1 and 2.2 subgroups by 54.1 g or 6.5% ( $P \leq 0.001$ ) and by 43.6 g - 5.3% ( $P \leq 0.01$ ) respectively.

On the basis of the research results, it can be concluded that of young pigs of Canadian breeding, the highest early ripeness and growth rate were for three-crossbred young pigs from 2.3 subgroup.

Comparing with young domestic breeding pigs from 1.1, 1.2 and 1.3 subgroups, pigs of Canadian breeding from 2.1, 2.2 and 2.3 subgroups reached live weight 24 and 22 days earlier at ( $P \leq 0.001$ )

Table 1 – Lifetime productivity of experimental young pigs ( $X \pm m_x$ ;  $n=20$ )

Group	Combination	Pre-slaughter live weight, kg	Age of reaching 100 kg mass, days	Average daily weight gain, g
1. Domestic breeding				
1.1	Largewhite	99.3±1.1	186±3	642.0±12.0
1.2	Largewhitex Landrace	101.2±0.9	182±2	672.1±13.4
1.3	Large whitex Landracex Duroc	102.3±1.0	178±2*	694.0±14.2**
2. Canadian breeding				
2.1	Yorkshirex Yorkshire	99.0±1.0	162±2.0	774.0±5.9
2.2	Yorkshirex Landrace	99.5±1.0	158±2.0	784.6±10.6**
2.3	Yorkshirex Landracex Duroc	102.0±1.1	156±2.0*	828.1±13.9***
<i>Note.</i> * $P \leq 0.05$ ; ** $P \leq 0.01$ ; *** $P \leq 0.001$ . Hereinafter, a dominance of young experimental subgroups over the control groups (1.1 and 2.1 subgroups) is indicated.				



respectively. In young stock of Canadian breeding from subgroups 2.1, 2.2 and 2.3 compared to subgroups 1.1, 1.2 and 1.3 of domestic breeding, the average daily weight gain was higher by 132.0 g or 15.9%, 112.5 g - 14.3% and 134.1 g - 16.2% respectively with ( $P \leq 0.001$ ).

Therefore, young Canadian breeding pigs significantly exceeded young stock of domestic breeding in average daily live weight gain.

The main indicator characterizing the meat productivity of pigs is the slaughter yield. The results of table 2 show that among young pigs of domestic breeding, the highest carcass mass was obtained from gilt of 1.3 subgroup - 69.6 kg, which is higher than in animals from 1.1 and 1.2 subgroups by 5.5 kg or 7.9% ( $P \leq 0.01$ ) and by 2.5 kg - 3.6%. The biggest mass of internal fat was obtained from young pigs of 1.1 subgroup - 3.2 kg, and the smallest - from gilt of 1.3 subgroup - 3.0 kg.

A higher slaughter yield was received from young pigs of 1.3 subgroup - 71.0%, which is higher compared to 1.1 and 1.2 subgroups by 3.2% ( $P \leq 0.05$ ) and 1.6%, respectively.

The slaughter results show that among the young domestic breeding pigs, the best slaughter indicators were shown by three-crossbred young stock.

Among young pigs of Canadian breeding, the highest carcass weight was obtained from 2.3 subgroup - 73.2 kg, which is higher compared to 2.1 and 2.2 subgroups by 4.6% ( $P \leq 0.05$ ) and 2.9% respectively. The highest mass of internal fat was obtained from pigs of 2.1 subgroup - 1.8 kg, and the lowest - from young stock of 2.1 subgroup - 1.6 kg. The highest results in slaughter yield were obtained from young pigs of 2.3 subgroup - 73.3%, which is higher compared to 2.1 and 2.2 subgroups of animals by 2.3% ( $P \leq 0.05$ ) and 1.2% respectively. So, the best slaughter rates among young pigs of Canadian breeding were obtained from three-breed crossbred young pigs of 2.3 subgroup.

Slaughter yield of young stock of Canadian breeding from 2.1, 2.2 and 2.3 subgroups compared to the young Canadian breeding of 1.1, 1.2 and 1.3 subgroups was higher by 3.2% ( $P \leq 0.05$ ), 2.7% and 2.3% respectively.

Table 2 – Slaughter indicators of young pigs ( $X \pm m_x$ ;  $n=10$ )

Group	Pre-slaughter live weight, kg	Carcass weight, kg	Mass of internal fat, kg	Deadweight, kg	Slaughter yield, %
1. Domestic breeding					
1.1	99.3±1.1	64.1±1.3	3.2±0.1	67.3±1.4	67.8±1.2
1.2	101.2±0.9	67.1±1.0	3.1±0.1	70.2±1.5	69.4±0.9
1.3	102.3±1.0	69.6±1.1**	3.0±0.1	72.6±1.2**	71.0±0.7*
2. Canadian breeding					
2.1	99.0±1.0	68.6±1.8	1.8±0.1	70.4±1.9	71.0±1.0
2.2	99.5±1.0	70.3±1.6	1.7±0.1	72.0±1.7	72.1±0.9
2.3	102.0±1.1	73.2±1.1*	1.6±0.1	74.8±1.1*	73.3±0.6*

Research results demonstrate that Canadian breeding is characterized by higher slaughter indicators. This certifies deeper and more effective breeding by fattening and meat qualities in pigs of Canadian breeding.

**Conclusion.** As a result of the research and production experiment, the biological features and productivity of domestic and Canadian breeding pigs were studied and it was established that:

- among domestic breeding young stock, the highest early ripeness and growth rate were for three-breed crossbred young pigs, obtained by crossing crossbred sows with Duroc boars;
- among Canadian breeding pigs, the highest precocity and growth rates were shown by three-breed crossbred stock obtained by mating of crossbred sows with Duroc boars;
- among the domestic breeding young pigs, the best slaughter indicators were shown by three-breed crossbred young stock;

- the best slaughter indicators among young pigs of Canadian breeding were for three-crossbred young stock;

- it was found that in comparison with the domestic breeding pigs, the three-breed crossbred young pigs reached the pre-slaughter live weight 22 days earlier; their average daily weight gains were higher by 134.1 g, and the slaughter yield was 2.3% higher. Thus, the three-breed commercial young stock of the Canadian breeding reliably surpasses the crossbred domestic breeding pigs.;

- a calculation of the economic efficiency of growing young pigs shows that the additional income from a realization of 1 pig of Canadian breeding, compared to domestic breeding, is 1.5 thousand rubles (\$ 23).

**Source of financing:** extrabudgetary funds of the Russian State Agrarian University - Moscow Agricultural Academy named after K.A. Timiryazev.

**С. А. Грикшас<sup>1</sup>, В. В. Калашников<sup>2</sup>, Г. К. Джанабекова<sup>3</sup>,  
Г. А. Фуников<sup>1</sup>, А. В. Овчинников<sup>1</sup>, Н. И. Кульмакова<sup>1</sup>, В. Т. Епифанов<sup>1</sup>,  
В. В. Храмов<sup>1</sup>, С. Н. Саримбекова<sup>3</sup>, М. Ш. Ережепова<sup>3</sup>**

<sup>1</sup>Жоғары білім беру саласындағы федералдық мемлекеттік бюджеттік жоғары білім беру мекемесі  
«Ресей мемлекеттік аграрлық университеті – К. А. Тимирязев атындағы

Мәскеу ауыл шаруашылық академиясы, Мәскеу, Ресей,

<sup>2</sup>Федералдық мемлекеттік бюджеттік мекеме «Бүкіл Ресейлік Жылқы Шаруашылығы»  
ғылыми-зерттеу институты, Рязан облысы Рыбинск ауданы Дивово ауылы, Ресей,

<sup>3</sup>Қазақ ұлттық аграрлық университет, Алматы, Қазақстан

### **КАНАДА ЖӘНЕ ОТАНДЫҚ СЕЛЕКЦИЯДАН ШЫҚҚАН ШОШҚАЛАРДЫҢ БИОЛОГИЯЛЫҚ ЖӘНЕ ӨНІМДІЛІК ЕРЕКШЕЛІКТЕРІ**

**Аннотация.** 2010-2015 жылдар аралығында шошқа өсірудің ірі кешендері жағдайында, шикізат аймағы ААҚ «Смолмясо» және ААҚ «Павловская Слобода» ет комбинатында зерттеумен өндірістік тәжірибе отандық және батыс селекциясы арқылы өсірілген шошқалардың өнімділігі мен биологиялық ерекшеліктерін зерттеу үшін жүргізілді. Тәжірибелік 2 айлық жас 100 кг тірі салмақ жинау үшін бақылау жағдайында өсірілді. Жас шошқалардың өсіру жағдайы үй, азықтандыру және сою шарттары бірдей болды. Жас шошқаның 95-105 кг тірі салмақ жинағаннан кейін олардың өнімділігі мен таза ет салмағының бағалауын өткіздік.

Зерттеу үрдісінде жас отандық және канадалық шошқалардың арасында ең жоғары өнімділік пен өсім қарқыны Дюрок тұқымының шошқасы мен будан шошқалармен будандастырылған үш тұқымдық жастармен сипатталды.

Бақылау сойысының нәтижелері жасотандық және канадалық шошқа арасында ең жақсы союыс көрсеткіш көрсеткен үштұқымдық будан жастары шошқалары.

Үштұқымдық будан жастары канадалық селекцияның отандық шошқадан ерекшелігі сойғанға дейін тірі салмағын 22 күнге ерте жинайды, орташа тірі салмақ жинау 134,1 г, сойыс шығысы 2,3% ға жоғары. Сондықтан, үш тұқымдық канадалық селекциядан шыққан тауарлық жас шошқалар отандық шошқалардан жоғары болып есептелді.

Жас өсірілген шошқалардың экономикалық тиімділігін есептеу Канаданың 1 басын сатудан түскен қосымша табыс отандық іріктеумен салыстырғанда 1,5 мың рубльді (23 АҚШ доллары) құрайды.

Зерттеу нәтижелері бойынша шошқа фермалары шошқа өсіру және шошқа етін өсірумен айналысады, шошқалардың бордақылау және ет сапасын арттыру үшін өсіруді күшейтеді. Бұрынғы бордақылау және ет сапасы жоғары сатылатын жас шошқаларды алу үшін канадалық шошқа шошқаларын кеңінен пайдалану ұсынылады.

**Түйін сөздер:** жас шошқа, отандық іріктеу, канадалық іріктеу, өсу қарқыны, күнделікті орташа өсу, сойыс салмағы, сойысшығысы.

С. А. Грикшас<sup>1</sup>, В. В. Калашников<sup>2</sup>, Г. К. Джанабекова<sup>3</sup>,  
Г. А. Фуников<sup>1</sup>, А. В. Овчинников<sup>1</sup>, Н. И. Кульмакова<sup>1</sup>, В. Т. Епифанов<sup>1</sup>,  
В. В. Храмцов<sup>1</sup>, С. Н. Саримбекова<sup>3</sup>, М. Ш. Ережепова<sup>3</sup>

<sup>1</sup>ФГБОУ ВО РГАУ-МСХА<sup>3</sup>Российский государственный аграрный университет –  
Московская сельскохозяйственная академия им. К.А. Тимирязева, Москва, Россия,

<sup>2</sup>Федеральное государственное бюджетное учреждение «Всероссийский научно-исследовательский  
институт коневодства», Рязанская область, Рыбновский район, п. Дивово, Россия,

<sup>3</sup>Казахский Национальный аграрный университет, Алматы, Казахстан

## ПРОДУКТИВНОСТЬ И БИОЛОГИЧЕСКИЕ ОСОБЕННОСТИ СВИНЕЙ ОТЕЧЕСТВЕННОЙ И КАНАДСКОЙ СЕЛЕКЦИИ

**Аннотация.** В условиях крупных свиноводческих комплексов в сырьевой зоне ОАО «Смолмясо» и ОАО «Мясокомбинат «Павловская Слобода» в течение 2010-2015 гг. был проведен научно-производственный опыт по изучению продуктивности и биологических особенностей свиней отечественной и западной селекции. Подопытный молодняк с 2-мес. возраста до достижения живой массы 100 кг содержался в условиях контрольного выращивания. Условия содержания, кормления и убоя молодняка свиней были идентичны. При достижении молодняком свиней живой массы в 95-105 кг провели оценку по собственной продуктивности и убойным показателям.

В процессе исследований было установлено, что среди молодняка отечественной и канадской селекций наиболее высокой скороспелостью и скоростью роста характеризовался трехпородный помесный молодняк, полученный при скрещивании помесных свиноматок с хряками породы дюрок. Результаты контрольного убоя показали, что среди молодняка отечественной и канадской селекций лучшими убойными показателями характеризовался трехпородный помесный молодняк.

Установлено, что трехпородный помесный молодняк канадской селекции по сравнению с отечественным достигал предубойной живой массы на 22 суток раньше, среднесуточные приросты живой массы были выше на 134,1 г, убойный выход был выше на 2,3%. Таким образом, трехпородный товарный молодняк канадской селекции по продуктивности достоверно превосходит свиней отечественной селекции.

Расчет экономической эффективности выращивания молодняка свиней показывает, что дополнительный доход от реализации 1 головы канадской селекции по сравнению с отечественной селекцией составляет 1,5 тыс. руб (23 доллара США).

На основе полученных результатов исследований было рекомендовано свинокомплексам, занимающимся разведением и выращиванием свиней отечественной селекции, усилить селекцию на повышение откормочных и мясных качеств свиней. Для получения товарного молодняка свиней с высокими откормочными и мясными качествами рекомендуется шире использовать свиней канадской селекции.

**Ключевые слова:** молодняк свиней, отечественная селекция, канадская селекция, скорость роста, среднесуточные приросты, убойная масса, убойный выход.

### Information about authors:

Griksas Styapas Antanovich, Doctor of Agricultural Sciences, Professor of the Department of Technology of Storage and Processing of Livestock Products, Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; [stepangr56@mail.ru](mailto:stepangr56@mail.ru); <https://orcid.org/0000-0002-7334-9719>

Kalashnikov Valery Vasilievich, Doctor of Agricultural Sciences, Professor, Academician of the Russian Academy of Sciences, Scientific Supervisor, All-Russian Scientific Research Institute of Horse Breeding, Divovo village, Rybnovsky District, Ryazan Region, Russia; [vniik08@mail.ru](mailto:vniik08@mail.ru); <https://orcid.org/0000-0003-2974-3732>

Dzhanabekova Gulmira Kumiskalieva, Doctor of Biological Sciences, Professor of the Department of Physiology, Morphology and Biochemistry named after Academician N. U. Bazanova, Kazakh National Agrarian University, Almaty, Kazakhstan; [guka.67@mail.ru](mailto:guka.67@mail.ru); <https://orcid.org/0000-0003-3940-729X>

Funikov Grigori Albertovich, Candidate of Agricultural Sciences, Executive Director of Viskotypak N. V. Company, Moscow, Russia; [gafvt@yandex.ru](mailto:gafvt@yandex.ru); <https://orcid.org/0000-0002-0471-3927>

Ovchinnikov Anatoly Viktorovich, Doctor of Agricultural Sciences, Professor of the Department of Private Animal Science, Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; [svinovod@rgau-msha.ru](mailto:svinovod@rgau-msha.ru); <https://orcid.org/0000-0002-7474-6345>

Kulmakova Natalia Ivanovna, Doctor of Agricultural Sciences, Professor of the Department of Morphology, Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; [kni11@mail.ru](mailto:kni11@mail.ru); <https://orcid.org/0000-0003-0372-6109>

Epifanov Victor Gennadiyevich, Doctor of Biological Sciences, Professor of the Department of Animal Feeding and Breeding, Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; [manager@nauchnik.org](mailto:manager@nauchnik.org); <https://orcid.org/0000-0003-1758-2492>

Khramtsov Vitaly Vasilyevich, Doctor of Agricultural Sciences, Professor of the Department of Veterinary and Morphology, Russian State Agrarian University – Moscow Agricultural Academy named after K. A. Timiryazev, Moscow, Russia; [komarovaoksana572@gmail.com](mailto:komarovaoksana572@gmail.com); <https://orcid.org/0000-0002-1674-2960>

Sarimbekova Saule Nurgaliyevna, Master of Science, Senior Teacher of the Department of Physiology, Morphology and Biochemistry named after Academician N. U. Bazanova, Kazakh National Agrarian University, Almaty, Kazakhstan; [sarimbekova@yandex.ru](mailto:sarimbekova@yandex.ru); <https://orcid.org/0000-0002-9625-5467>

Erezhepova Makpal Shamakhanovna, Master of Science, Assistant Professor of the Department of Physiology, Morphology and Biochemistry named after Academician N. U. Bazanova, Kazakh National Agrarian University, Almaty, Kazakhstan; [princessa.maki@mail.ru](mailto:princessa.maki@mail.ru); <https://orcid.org/0000-0002-3945-4611>

## REFERENCES

- [1] Belousov N. (2016). Effective development of pig husbandry. Pig production. Moscow. N 2. P. 66-67 (in Russ.).
- [2] Grikshas S.A. (2011). Ways and methods of improving the breeding and productive qualities of specialized lines and types of pigs and the effectiveness of use in hybridization. Monograph. Moscow: MSAU. 116 p. (in Russ.).
- [3] Suslina E.N. (2017). State and development of the breeding sector of the domestic pig husbandry. Pig production. Moscow. N 4. P. 4-6 (in Russ.).
- [4] Russia ranks 5th in the world in pork production [Electronic resource]. (2016). Electron. text data. – Access mode: <http://www.servis-expo.ru/news/rossiya-vyshla-na-5-mesto-v-mire-po-proizvodstvu-svininy>. Date of applying: 29.12.2016.
- [5] Tatulov Yu.V., Kolomiets N.N., Rozanov A.V., Grikshas S.A. (2001). To the problem of creating industrially suitable genotypes of pigs // Meat industry. N 1. P. 40-42 (in Russ.).
- [6] Chernukha I.M., Tatulov Yu.V., Veselov P.P., Sus I.V., Grikshas S.A., Kolomiets N.N. (2004). Recommendations for the use of industrially suitable pig genotypes in the meat industry. Moscow: MSHA publishing house, 2004. P. 8 (in Russ.).
- [7] Velichko V.A., Komlatsky V.I., Velichko L.F. (2017). The path to import substitution - the use of foreign breeding pigs. Scientific journal KubGAU. Kuban, 2017. N 130(06). P. 1-8 (in Russ.).
- [8] Pokhodnya G.S., Ivchenko A.N., Korobov D.V., Malakhova T.A. (2016). The effectiveness of cross-breeding of large white sows with boars of Duroc breed of import breeding // Tavrichesky scientific reviewer. Crimea. N 5(10). May. P. 172-178 (in Russ.).
- [9] Solovyh A., Ovchinnikov A., Khrenova O. (2005) The best maternal breed - Large white. Animal husbandry of Russia. Moscow. N 12. P. 22-23 (in Russ.).
- [10] Sokolov N.V., Zelkova N.G. (2018). Comparative productivity of pigs of large white and landrace breeds at linear breeding and crossing. Achievements of science and technology of agriculture. Moscow. Vol. 34, N 4. P. 54-58. DOI: 10.24411//0235-2451-2018-10413 (in Russ.).
- [11] Burtseva S.V., Pautova L.N. (2016). The influence of the Yorkshire blood on meat quality of pigs // Bulletin of the Altai State University. Altai. N 10(144). P. 102-106 (in Russ.).
- [12] Gubanova N.S. (2013). Biological and productive features of Canadian breeding pigs: Author's abstract ... dis. can. biol. sciences. Moscow. 20 p. (in Russ.).
- [13] Vodyannikov V.I., Shkalenko V.V., Ruzhenikov F.V., Zemlyakov R.N. (2010). Productivity and quality of Canadian pork in the Lower Volga region. Pig husbandry. Moscow. N 6. P. 14-15 (in Russ.).
- [14] Bekenev V.A., Frolova V.I., Botsan I.V., Frolova Yu.V., Kharseeva M.I., Zabolotnaya A.A., Gaptar S.I., Golovko A.N. (2012). The results of the use of imported meat breeds of pigs when crossed in Siberia. Achievements of science and technology of agriculture. Moscow. N 7. P. 67-69 (in Russ.).
- [15] Funikov G.A. (2001). The productivity and quality of the meat of large white breed pigs in purebred breeding and crossing with boars of large black, Landrace and Duroc breeds: Abstract ... dis. can. agricultural sciences. 17 p. (in Russ.).
- [16] Takahashi K. (1996). Structural weakening of skeletal muscle tissue during post-mortem aging of meat: the non-enzymatic mechanism of meat tenderization. Meat science. 43. P. 67-S80.
- [17] Fredeen X.T. (1969). Breed structure and population dynamics of the Canadian Yorkshire pig: Canadian Journal of Animal Science. 49(3): 291-304 (doi: 10.4141/cjas69-040).
- [18] Norms and diets of feeding farm animals. (2003). Reference manual. 3rd edition revised and enlarged / Ed. A.P. Kalashnikov, V.I. Fisinin, V.V. Shcheglova, N.I. Kleimenova. Moscow. 456 p. (in Russ.).
- [19] Methodological recommendations of the Academy of Agricultural Sciences on the assessment of meat productivity, meat quality and subcutaneous fat of pigs. (1978). Moscow: Agricultural Sciences. 43 p. (in Russ.).
- [20] Gataulin A.M. (1992). The system of applied statistical and mathematical methods for processing experimental data in agriculture (ISSN 5-7230-0001-2). 2 parts. Moscow. Ed. TSHA. 23.5 p.p.
- [21] Baimukanov D.A., Pristupa V.N., Kolosov Yu.A., Donnik I.M., Torosyan D.S., Kolosov A.Yu., Orlova O.N., Yuldashbayev Yu.A., Chylbak-ool S.O. (2019). Improvement of breeding and productive traits of Kalmyk cattle breed // Bulletin of National academy of sciences of the Republic of Kazakhstan. Vol. 1, N 378 (2019). P. 128-145. ISSN 2518-1467 (Online), ISSN 1991-3494 (Print). <https://doi.org/10.32014/2019.2518-1467.51>

---

---

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

**ISSN 2518-1467 (Online), ISSN 1991-3494 (Print)**

<http://www.bulletin-science.kz/index.php/en/>

Редакторы *М. С. Ахметова, Т. М. Апендиев, Д. С. Аленов*  
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 10.10.2019.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
13,9 п.л. Тираж 500. Заказ 5.